

## 9.8 Power Series

- Understand the definition of a power series.
- Find the radius and interval of convergence of a power series.
- Determine the endpoint convergence of a power series.
- Differentiate and integrate a power series.

### Power Series

In Section 9.7, you were introduced to the concept of approximating functions by Taylor polynomials. For instance, the function  $f(x) = e^x$  can be *approximated* by its Maclaurin polynomials as follows.

$$e^x \approx 1 + x \quad \text{1st-degree polynomial}$$

$$e^x \approx 1 + x + \frac{x^2}{2!} \quad \text{2nd-degree polynomial}$$

$$e^x \approx 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} \quad \text{3rd-degree polynomial}$$

$$e^x \approx 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} \quad \text{4th-degree polynomial}$$

$$e^x \approx 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} \quad \text{5th-degree polynomial}$$

In that section, you saw that the higher the degree of the approximating polynomial, the better the approximation becomes.

In this and the next two sections, you will see that several important types of functions, including

$$f(x) = e^x$$

can be represented *exactly* by an infinite series called a **power series**. For example, the power series representation for  $e^x$  is

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

For each real number  $x$ , it can be shown that the infinite series on the right converges to the number  $e^x$ . Before doing this, however, some preliminary results dealing with power series will be discussed—beginning with the following definition.

#### DEFINITION OF POWER SERIES

If  $x$  is a variable, then an infinite series of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots + a_n x^n + \cdots$$

is called a **power series**. More generally, an infinite series of the form

$$\sum_{n=0}^{\infty} a_n (x - c)^n = a_0 + a_1 (x - c) + a_2 (x - c)^2 + \cdots + a_n (x - c)^n + \cdots$$

is called a **power series centered at  $c$** , where  $c$  is a constant.

**NOTE** To simplify the notation for power series, we agree that  $(x - c)^0 = 1$ , even if  $x = c$ .

#### EXPLORATION

**Graphical Reasoning** Use a graphing utility to approximate the graph of each power series near  $x = 0$ . (Use the first several terms of each series.) Each series represents a well-known function. What is the function?

a.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{n!}$

b.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$

c.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$

d.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$

e.  $\sum_{n=0}^{\infty} \frac{2^n x^n}{n!}$

**EXAMPLE 1** Power Series

a. The following power series is centered at 0.

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \cdots$$

b. The following power series is centered at  $-1$ .

$$\sum_{n=0}^{\infty} (-1)^n (x + 1)^n = 1 - (x + 1) + (x + 1)^2 - (x + 1)^3 + \cdots$$

c. The following power series is centered at 1.

$$\sum_{n=1}^{\infty} \frac{1}{n} (x - 1)^n = (x - 1) + \frac{1}{2} (x - 1)^2 + \frac{1}{3} (x - 1)^3 + \cdots$$

**Radius and Interval of Convergence**


A power series in  $x$  can be viewed as a function of  $x$

$$f(x) = \sum_{n=0}^{\infty} a_n (x - c)^n$$

where the *domain of  $f$*  is the set of all  $x$  for which the power series converges. Determination of the domain of a power series is the primary concern in this section. Of course, every power series converges at its center  $c$  because

$$\begin{aligned} f(c) &= \sum_{n=0}^{\infty} a_n (c - c)^n \\ &= a_0(1) + 0 + 0 + \cdots + 0 + \cdots \\ &= a_0. \end{aligned}$$

So,  $c$  always lies in the domain of  $f$ . The following important theorem states that the domain of a power series can take three basic forms: a single point, an interval centered at  $c$ , or the entire real line, as shown in Figure 9.17. A proof is given in Appendix A.



The domain of a power series has only three basic forms: a single point, an interval centered at  $c$ , or the entire real line.

**Figure 9.17**

**THEOREM 9.20 CONVERGENCE OF A POWER SERIES**

For a power series centered at  $c$ , precisely one of the following is true.

1. The series converges only at  $c$ .
2. There exists a real number  $R > 0$  such that the series converges absolutely for  $|x - c| < R$ , and diverges for  $|x - c| > R$ .
3. The series converges absolutely for all  $x$ .

The number  $R$  is the **radius of convergence** of the power series. If the series converges only at  $c$ , the radius of convergence is  $R = 0$ , and if the series converges for all  $x$ , the radius of convergence is  $R = \infty$ . The set of all values of  $x$  for which the power series converges is the **interval of convergence** of the power series.

**STUDY TIP** To determine the radius of convergence of a power series, use the Ratio Test, as demonstrated in Examples 2, 3, and 4.

### EXAMPLE 2 Finding the Radius of Convergence

Find the radius of convergence of  $\sum_{n=0}^{\infty} n!x^n$ .

**Solution** For  $x = 0$ , you obtain

$$f(0) = \sum_{n=0}^{\infty} n!0^n = 1 + 0 + 0 + \cdots = 1.$$

For any fixed value of  $x$  such that  $|x| > 0$ , let  $u_n = n!x^n$ . Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| \\ &= |x| \lim_{n \rightarrow \infty} (n+1) \\ &= \infty. \end{aligned}$$

Therefore, by the Ratio Test, the series diverges for  $|x| > 0$  and converges only at its center, 0. So, the radius of convergence is  $R = 0$ .

### EXAMPLE 3 Finding the Radius of Convergence

Find the radius of convergence of

$$\sum_{n=0}^{\infty} 3(x-2)^n.$$

**Solution** For  $x \neq 2$ , let  $u_n = 3(x-2)^n$ . Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{3(x-2)^{n+1}}{3(x-2)^n} \right| \\ &= \lim_{n \rightarrow \infty} |x-2| \\ &= |x-2|. \end{aligned}$$

By the Ratio Test, the series converges if  $|x-2| < 1$  and diverges if  $|x-2| > 1$ . Therefore, the radius of convergence of the series is  $R = 1$ .

### EXAMPLE 4 Finding the Radius of Convergence

Find the radius of convergence of

$$\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}.$$


**Solution** Let  $u_n = (-1)^n x^{2n+1}/(2n+1)!$ . Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1} x^{2n+3}}{(2n+3)!}}{\frac{(-1)^n x^{2n+1}}{(2n+1)!}} \right| \\ &= \lim_{n \rightarrow \infty} \frac{x^2}{(2n+3)(2n+2)}. \end{aligned}$$

For any fixed value of  $x$ , this limit is 0. So, by the Ratio Test, the series converges for all  $x$ . Therefore, the radius of convergence is  $R = \infty$ . ■

## Endpoint Convergence

Note that for a power series whose radius of convergence is a finite number  $R$ , Theorem 9.20 says nothing about the convergence at the *endpoints* of the interval of convergence. Each endpoint must be tested separately for convergence or divergence. As a result, the interval of convergence of a power series can take any one of the six forms shown in Figure 9.18.



Intervals of convergence

Figure 9.18

### EXAMPLE 5 Finding the Interval of Convergence

Find the interval of convergence of  $\sum_{n=1}^{\infty} \frac{x^n}{n}$ .

**Solution** Letting  $u_n = x^n/n$  produces

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{x^{n+1}}{n+1}}{\frac{x^n}{n}} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{nx}{n+1} \right| \\ &= |x|. \end{aligned}$$

So, by the Ratio Test, the radius of convergence is  $R = 1$ . Moreover, because the series is centered at 0, it converges in the interval  $(-1, 1)$ . This interval, however, is not necessarily the *interval of convergence*. To determine this, you must test for convergence at each endpoint. When  $x = 1$ , you obtain the *divergent* harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n} = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \cdots \quad \text{Diverges when } x = 1$$

When  $x = -1$ , you obtain the *convergent* alternating harmonic series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n} = -1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \cdots \quad \text{Converges when } x = -1$$

So, the interval of convergence for the series is  $[-1, 1)$ , as shown in Figure 9.19.




Figure 9.19

**EXAMPLE 6** Finding the Interval of Convergence

Find the interval of convergence of  $\sum_{n=0}^{\infty} \frac{(-1)^n(x+1)^n}{2^n}$ .

**Solution** Letting  $u_n = (-1)^n(x+1)^n/2^n$  produces

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1}(x+1)^{n+1}}{2^{n+1}}}{\frac{(-1)^n(x+1)^n}{2^n}} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{2^n(x+1)}{2^{n+1}} \right| \\ &= \left| \frac{x+1}{2} \right|. \end{aligned}$$

By the Ratio Test, the series converges if  $|(x+1)/2| < 1$  or  $|x+1| < 2$ . So, the radius of convergence is  $R = 2$ . Because the series is centered at  $x = -1$ , it will converge in the interval  $(-3, 1)$ . Furthermore, at the endpoints you have

$$\sum_{n=0}^{\infty} \frac{(-1)^n(-2)^n}{2^n} = \sum_{n=0}^{\infty} \frac{2^n}{2^n} = \sum_{n=0}^{\infty} 1 \quad \text{Diverges when } x = -3$$

and

$$\sum_{n=0}^{\infty} \frac{(-1)^n(2)^n}{2^n} = \sum_{n=0}^{\infty} (-1)^n \quad \text{Diverges when } x = 1$$

both of which diverge. So, the interval of convergence is  $(-3, 1)$ , as shown in Figure 9.20.

Interval:  $(-3, 1)$   
Radius:  $R = 2$




Figure 9.20

**EXAMPLE 7** Finding the Interval of Convergence

Find the interval of convergence of

$$\sum_{n=1}^{\infty} \frac{x^n}{n^2}$$

**Solution** Letting  $u_n = x^n/n^2$  produces

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}/(n+1)^2}{x^n/n^2} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{n^2x}{(n+1)^2} \right| = |x|. \end{aligned}$$


So, the radius of convergence is  $R = 1$ . Because the series is centered at  $x = 0$ , it converges in the interval  $(-1, 1)$ . When  $x = 1$ , you obtain the convergent  $p$ -series

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots \quad \text{Converges when } x = 1$$

When  $x = -1$ , you obtain the convergent alternating series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{1}{1^2} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} - \cdots \quad \text{Converges when } x = -1$$

Therefore, the interval of convergence for the given series is  $[-1, 1]$ . ■



The Granger Collection

**JAMES GREGORY (1638–1675)**

One of the earliest mathematicians to work with power series was a Scotsman, James Gregory. He developed a power series method for interpolating table values—a method that was later used by Brook Taylor in the development of Taylor polynomials and Taylor series.

## Differentiation and Integration of Power Series

Power series representation of functions has played an important role in the development of calculus. In fact, much of Newton's work with differentiation and integration was done in the context of power series—especially his work with complicated algebraic functions and transcendental functions. Euler, Lagrange, Leibniz, and the Bernoullis all used power series extensively in calculus.

Once you have defined a function with a power series, it is natural to wonder how you can determine the characteristics of the function. Is it continuous? Differentiable? Theorem 9.21, which is stated without proof, answers these questions.

### THEOREM 9.21 PROPERTIES OF FUNCTIONS DEFINED BY POWER SERIES

If the function given by

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} a_n(x-c)^n \\ &= a_0 + a_1(x-c) + a_2(x-c)^2 + a_3(x-c)^3 + \cdots \end{aligned}$$

has a radius of convergence of  $R > 0$ , then, on the interval  $(c - R, c + R)$ ,  $f$  is differentiable (and therefore continuous). Moreover, the derivative and antiderivative of  $f$  are as follows.

- $$\begin{aligned} f'(x) &= \sum_{n=1}^{\infty} n a_n(x-c)^{n-1} \\ &= a_1 + 2a_2(x-c) + 3a_3(x-c)^2 + \cdots \end{aligned}$$
- $$\begin{aligned} \int f(x) dx &= C + \sum_{n=0}^{\infty} a_n \frac{(x-c)^{n+1}}{n+1} \\ &= C + a_0(x-c) + a_1 \frac{(x-c)^2}{2} + a_2 \frac{(x-c)^3}{3} + \cdots \end{aligned}$$

The *radius of convergence* of the series obtained by differentiating or integrating a power series is the same as that of the original power series. The *interval of convergence*, however, may differ as a result of the behavior at the endpoints.

Theorem 9.21 states that, in many ways, a function defined by a power series behaves like a polynomial. It is continuous in its interval of convergence, and both its derivative and its antiderivative can be determined by differentiating and integrating each term of the given power series. For instance, the derivative of the power series

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} \frac{x^n}{n!} \\ &= 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots \end{aligned}$$

is

$$\begin{aligned} f'(x) &= 1 + (2) \frac{x}{2} + (3) \frac{x^2}{3!} + (4) \frac{x^3}{4!} + \cdots \\ &= 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots \\ &= f(x). \end{aligned}$$

Notice that  $f'(x) = f(x)$ . Do you recognize this function?

**EXAMPLE 8** Intervals of Convergence for  $f(x)$ ,  $f'(x)$ , and  $\int f(x) dx$

Consider the function given by

$$f(x) = \sum_{n=1}^{\infty} \frac{x^n}{n} = x + \frac{x^2}{2} + \frac{x^3}{3} + \dots$$

Find the interval of convergence for each of the following.

- a.  $\int f(x) dx$       b.  $f(x)$       c.  $f'(x)$

**Solution** By Theorem 9.21, you have

$$\begin{aligned} f'(x) &= \sum_{n=1}^{\infty} x^{n-1} \\ &= 1 + x + x^2 + x^3 + \dots \end{aligned}$$

and

$$\begin{aligned} \int f(x) dx &= C + \sum_{n=1}^{\infty} \frac{x^{n+1}}{n(n+1)} \\ &= C + \frac{x^2}{1 \cdot 2} + \frac{x^3}{2 \cdot 3} + \frac{x^4}{3 \cdot 4} + \dots \end{aligned}$$

By the Ratio Test, you can show that each series has a radius of convergence of  $R = 1$ . Considering the interval  $(-1, 1)$ , you have the following.

- a. For  $\int f(x) dx$ , the series

$$\sum_{n=1}^{\infty} \frac{x^{n+1}}{n(n+1)} \quad \text{Interval of convergence: } [-1, 1]$$

converges for  $x = \pm 1$ , and its interval of convergence is  $[-1, 1]$ . See Figure 9.21(a).

- b. For  $f(x)$ , the series


$$\sum_{n=1}^{\infty} \frac{x^n}{n} \quad \text{Interval of convergence: } [-1, 1)$$

converges for  $x = -1$  and diverges for  $x = 1$ . So, its interval of convergence is  $[-1, 1)$ . See Figure 9.21(b).

- c. For  $f'(x)$ , the series

$$\sum_{n=1}^{\infty} x^{n-1} \quad \text{Interval of convergence: } (-1, 1)$$

diverges for  $x = \pm 1$ , and its interval of convergence is  $(-1, 1)$ . See Figure 9.21(c).



**Figure 9.21**

From Example 8, it appears that of the three series, the one for the derivative,  $f'(x)$ , is the least likely to converge at the endpoints. In fact, it can be shown that if the series for  $f'(x)$  converges at the endpoints  $x = c \pm R$ , the series for  $f(x)$  will also converge there.

## 9.8

## Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–4, state where the power series is centered.

1.  $\sum_{n=0}^{\infty} nx^n$

2.  $\sum_{n=1}^{\infty} \frac{(-1)^n 1 \cdot 3 \cdot \dots \cdot (2n-1)}{2^n n!} x^n$

3.  $\sum_{n=1}^{\infty} \frac{(x-2)^n}{n^3}$

4.  $\sum_{n=0}^{\infty} \frac{(-1)^n (x-\pi)^{2n}}{(2n)!}$

In Exercises 5–10, find the radius of convergence of the power series.

5.  $\sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n+1}$

6.  $\sum_{n=0}^{\infty} (4x)^n$

7.  $\sum_{n=1}^{\infty} \frac{(4x)^n}{n^2}$

8.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{5^n}$

9.  $\sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}$

10.  $\sum_{n=0}^{\infty} \frac{(2n)! x^{2n}}{n!}$

In Exercises 11–34, find the interval of convergence of the power series. (Be sure to include a check for convergence at the endpoints of the interval.)

11.  $\sum_{n=0}^{\infty} \left(\frac{x}{4}\right)^n$

12.  $\sum_{n=0}^{\infty} \left(\frac{x}{7}\right)^n$

13.  $\sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n}$

14.  $\sum_{n=0}^{\infty} (-1)^{n+1} (n+1)x^n$

15.  $\sum_{n=0}^{\infty} \frac{x^{5n}}{n!}$

16.  $\sum_{n=0}^{\infty} \frac{(3x)^n}{(2n)!}$

17.  $\sum_{n=0}^{\infty} (2n)! \left(\frac{x}{3}\right)^n$

18.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{(n+1)(n+2)}$

19.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{4^n}$

20.  $\sum_{n=0}^{\infty} \frac{(-1)^n n!(x-5)^n}{3^n}$

21.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-4)^n}{n9^n}$

22.  $\sum_{n=0}^{\infty} \frac{(x-3)^{n+1}}{(n+1)4^{n+1}}$

23.  $\sum_{n=0}^{\infty} \frac{(-1)^{n+1} (x-1)^{n+1}}{n+1}$

24.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-2)^n}{n2^n}$

25.  $\sum_{n=1}^{\infty} \frac{(x-3)^{n-1}}{3^{n-1}}$

26.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$

27.  $\sum_{n=1}^{\infty} \frac{n}{n+1} (-2x)^{n-1}$

28.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{n!}$

29.  $\sum_{n=0}^{\infty} \frac{x^{3n+1}}{(3n+1)!}$

30.  $\sum_{n=1}^{\infty} \frac{n! x^n}{(2n)!}$

31.  $\sum_{n=1}^{\infty} \frac{2 \cdot 3 \cdot 4 \cdot \dots \cdot (n+1)x^n}{n!}$

32.  $\sum_{n=1}^{\infty} \left[ \frac{2 \cdot 4 \cdot 6 \cdot \dots \cdot 2n}{3 \cdot 5 \cdot 7 \cdot \dots \cdot (2n+1)} \right] x^{2n+1}$

33.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} 3 \cdot 7 \cdot 11 \cdot \dots \cdot (4n-1)(x-3)^n}{4^n}$

34.  $\sum_{n=1}^{\infty} \frac{n!(x+1)^n}{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}$

In Exercises 35 and 36, find the radius of convergence of the power series, where  $c > 0$  and  $k$  is a positive integer.

35.  $\sum_{n=1}^{\infty} \frac{(x-c)^{n-1}}{c^{n-1}}$

36.  $\sum_{n=0}^{\infty} \frac{(n!)^k x^n}{(kn)!}$

In Exercises 37–40, find the interval of convergence of the power series. (Be sure to include a check for convergence at the endpoints of the interval.)

37.  $\sum_{n=0}^{\infty} \left(\frac{x}{k}\right)^n, k > 0$

38.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-c)^n}{nc^n}$

39.  $\sum_{n=1}^{\infty} \frac{k(k+1)(k+2) \cdot \dots \cdot (k+n-1)x^n}{n!}, k \geq 1$

40.  $\sum_{n=1}^{\infty} \frac{n!(x-c)^n}{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}$

In Exercises 41–44, write an equivalent series with the index of summation beginning at  $n = 1$ .

41.  $\sum_{n=0}^{\infty} \frac{x^n}{n!}$

42.  $\sum_{n=0}^{\infty} (-1)^{n+1} (n+1)x^n$

43.  $\sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$

44.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$

In Exercises 45–48, find the intervals of convergence of (a)  $f(x)$ , (b)  $f'(x)$ , (c)  $f''(x)$ , and (d)  $\int f(x) dx$ . Include a check for convergence at the endpoints of the interval.

45.  $f(x) = \sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n$

46.  $f(x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-5)^n}{n5^n}$



47.  $f(x) = \sum_{n=0}^{\infty} \frac{(-1)^{n+1} (x-1)^{n+1}}{n+1}$

48.  $f(x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-2)^n}{n}$



**Writing** In Exercises 49–52, match the graph of the first 10 terms of the sequence of partial sums of the series

$$g(x) = \sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n$$

with the indicated value of the function. [The graphs are labeled (a), (b), (c), and (d).] Explain how you made your choice.







49.  $g(1)$

50.  $g(2)$





51.  $g(3.1)$

52.  $g(-2)$

**Writing** In Exercises 53–56, match the graph of the first 10 terms of the sequence of partial sums of the series

$$g(x) = \sum_{n=0}^{\infty} (2x)^n$$

with the indicated value of the function. [The graphs are labeled (a), (b), (c), and (d).] Explain how you made your choice.



53.  $g\left(\frac{1}{8}\right)$

54.  $g\left(-\frac{1}{8}\right)$

55.  $g\left(\frac{9}{16}\right)$

56.  $g\left(\frac{3}{8}\right)$

**WRITING ABOUT CONCEPTS**

- 57. Define a power series centered at  $c$ .
- 58. Describe the radius of convergence of a power series. Describe the interval of convergence of a power series.
- 59. Describe the three basic forms of the domain of a power series.
- 60. Describe how to differentiate and integrate a power series with a radius of convergence  $R$ . Will the series resulting from the operations of differentiation and integration have a different radius of convergence? Explain.

**WRITING ABOUT CONCEPTS (continued)**

- 61. Give examples that show that the convergence of a power series at an endpoint of its interval of convergence may be either conditional or absolute. Explain your reasoning.

**CAPSTONE**

- 62. Write a power series that has the indicated interval of convergence. Explain your reasoning.
  - (a)  $(-2, 2)$
  - (b)  $(-1, 1]$
  - (c)  $(-1, 0)$
  - (d)  $[-2, 6)$

63. Let  $f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$  and  $g(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$ .

- (a) Find the intervals of convergence of  $f$  and  $g$ .
- (b) Show that  $f'(x) = g(x)$ .
- (c) Show that  $g'(x) = -f(x)$ .
- (d) Identify the functions  $f$  and  $g$ .

64. Let  $f(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ .

- (a) Find the interval of convergence of  $f$ .
- (b) Show that  $f'(x) = f(x)$ .
- (c) Show that  $f(0) = 1$ .
- (d) Identify the function  $f$ .

In Exercises 65–70, show that the function represented by the power series is a solution of the differential equation.

65.  $y = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$ ,  $y'' + y = 0$

66.  $y = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$ ,  $y'' + y = 0$

67.  $y = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$ ,  $y'' - y = 0$

68.  $y = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}$ ,  $y'' - y = 0$

69.  $y = \sum_{n=0}^{\infty} \frac{x^{2n}}{2^n n!}$ ,  $y'' - xy' - y = 0$

70.  $y = 1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{4n}}{2^{2n} n! \cdot 3 \cdot 7 \cdot 11 \cdots (4n-1)}$ ,  $y'' + x^2 y = 0$

71. **Bessel Function** The Bessel function of order 0 is

$$J_0(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{2^{2k} (k!)^2}$$

- (a) Show that the series converges for all  $x$ .
- (b) Show that the series is a solution of the differential equation  $x^2 J_0'' + x J_0' + x^2 J_0 = 0$ .
- (c) Use a graphing utility to graph the polynomial composed of the first four terms of  $J_0$ .
- (d) Approximate  $\int_0^1 J_0 dx$  accurate to two decimal places.

72. **Bessel Function** The Bessel function of order 1 is

$$J_1(x) = x \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{2^{2k+1} k!(k+1)!}$$

- (a) Show that the series converges for all  $x$ .  
 (b) Show that the series is a solution of the differential equation  $x^2 J_1'' + x J_1' + (x^2 - 1) J_1 = 0$ .  
 (c) Use a graphing utility to graph the polynomial composed of the first four terms of  $J_1$ .  
 (d) Show that  $J_0'(x) = -J_1(x)$ .

**CAS** In Exercises 73–76, the series represents a well-known function. Use a computer algebra system to graph the partial sum  $S_{10}$  and identify the function from the graph.

73.  $f(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$       74.  $f(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$

75.  $f(x) = \sum_{n=0}^{\infty} (-1)^n x^n, \quad -1 < x < 1$

76.  $f(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}, \quad -1 \leq x \leq 1$

**77. Investigation** The interval of convergence of the geometric series  $\sum_{n=0}^{\infty} \left(\frac{x}{4}\right)^n$  is  $(-4, 4)$ .

- (a) Find the sum of the series when  $x = \frac{5}{2}$ . Use a graphing utility to graph the first six terms of the sequence of partial sums and the horizontal line representing the sum of the series.  
 (b) Repeat part (a) for  $x = -\frac{5}{2}$ .  
 (c) Write a short paragraph comparing the rates of convergence of the partial sums with the sums of the series in parts (a) and (b). How do the plots of the partial sums differ as they converge toward the sum of the series?  
 (d) Given any positive real number  $M$ , there exists a positive integer  $N$  such that the partial sum

$$\sum_{n=0}^N \left(\frac{5}{4}\right)^n > M.$$

Use a graphing utility to complete the table.

$M$	10	100	1000	10,000
$N$				

**78. Investigation** The interval of convergence of the series  $\sum_{n=0}^{\infty} (3x)^n$  is  $(-\frac{1}{3}, \frac{1}{3})$ .

- (a) Find the sum of the series when  $x = \frac{1}{6}$ . Use a graphing utility to graph the first six terms of the sequence of partial sums and the horizontal line representing the sum of the series.  
 (b) Repeat part (a) for  $x = -\frac{1}{6}$ .  
 (c) Write a short paragraph comparing the rates of convergence of the partial sums with the sums of the series in parts (a) and (b). How do the plots of the partial sums differ as they converge toward the sum of the series?

(d) Given any positive real number  $M$ , there exists a positive integer  $N$  such that the partial sum

$$\sum_{n=0}^N \left(3 \cdot \frac{2}{3}\right)^n > M.$$

Use a graphing utility to complete the table.

$M$	10	100	1000	10,000
$N$				

**True or False?** In Exercises 79–82, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

79. If the power series  $\sum_{n=1}^{\infty} a_n x^n$  converges for  $x = 2$ , then it also converges for  $x = -2$ .  
 80. It is possible to find a power series whose interval of convergence is  $[0, \infty)$ .  
 81. If the interval of convergence for  $\sum_{n=0}^{\infty} a_n x^n$  is  $(-1, 1)$ , then the interval of convergence for  $\sum_{n=0}^{\infty} a_n (x-1)^n$  is  $(0, 2)$ .

82. If  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  converges for  $|x| < 2$ , then

$$\int_0^1 f(x) dx = \sum_{n=0}^{\infty} \frac{a_n}{n+1}.$$

83. Prove that the power series

$$\sum_{n=0}^{\infty} \frac{(n+p)!}{n!(n+q)!} x^n$$

has a radius of convergence of  $R = \infty$  if  $p$  and  $q$  are positive integers.

84. Let  $g(x) = 1 + 2x + x^2 + 2x^3 + x^4 + \cdots$ , where the coefficients are  $c_{2n} = 1$  and  $c_{2n+1} = 2$  for  $n \geq 0$ .

- (a) Find the interval of convergence of the series.  
 (b) Find an explicit formula for  $g(x)$ .

85. Let  $f(x) = \sum_{n=0}^{\infty} c_n x^n$ , where  $c_{n+3} = c_n$  for  $n \geq 0$ .

- (a) Find the interval of convergence of the series.  
 (b) Find an explicit formula for  $f(x)$ .

86. Prove that if the power series  $\sum_{n=0}^{\infty} c_n x^n$  has a radius of convergence of  $R$ , then  $\sum_{n=0}^{\infty} c_n x^{2n}$  has a radius of convergence of  $\sqrt{R}$ .

87. For  $n > 0$ , let  $R > 0$  and  $c_n > 0$ . Prove that if the interval of convergence of the series  $\sum_{n=0}^{\infty} c_n (x-x_0)^n$  is  $[x_0 - R, x_0 + R]$  then the series converges conditionally at  $x_0 - R$ .